

A L^AT_EX Tutorial

David Wagner

U.C. Berkeley

daw@cs.berkeley.edu

Why L^AT_EX?

- Best way to typeset mathematics
(vastly better than MS-Word)
- Most popular typesetting system among computer scientists
- Can be used to produce beautiful output

How to use L^AT_EX?

1. Create `hw1.tex` with your favorite text editor.
2. Compile: `latex hw1.tex`
This produces `hw1.dvi`.
3. Make PDF: `dvipdf hw1.dvi hw1.pdf`
This produces `hw1.pdf`.
4. View: `acroread hw1.pdf`

Sample document

Input:

```
\documentclass{article}  
\begin{document}  
Hello, world!  
\end{document}
```

Output:

```
Hello, world!
```

Sample document

From here on in, I'll omit the `\documentclass{...}`,
`\begin{document}` and `\end{document}`. Thus:

Input: `Hello, world!`

Output: `Hello, world!`

Paragraphs

```
Line breaks in latex source  
are irrelevant, except that  
blank lines indicate the start  
of a new paragraph.
```

```
See how this  
works?
```

Line breaks in latex source are irrelevant, except that blank lines indicate the start of a new paragraph.

See how this works?

Commands

Commands use a backslash.

Arguments are indicated with curly braces.

Let's try some `\emph{italized}` text.

Commands use a backslash. Arguments are indicated with curly braces.

Let's try some *italized* text.

Mathematics

In-line mathematics is enclosed within dollar signs.

$\$1+2=3\$$ looks better than $1+2=3$.

$\$12x+5 > y\$$ looks better than $12x+5 > y$.

In-line mathematics is enclosed within dollar signs.

$1 + 2 = 3$ looks better than $1+2=3$.

$12x + 5 > y$ looks better than $12x+5 > y$.

Superscripts

Squaring: x^2 . Higher powers: x^n .

Or: $(2n+1)^3 = 8n^3 + 12n^2 + 6n + 1$.

Warning: If you have more than one character in the exponent, you must use curly braces for grouping.

Correct: x^{2n} . Wrong: x^2n .

Squaring: x^2 . Higher powers: x^n . Or: $(2n+1)^3 = 8n^3 + 12n^2 + 6n + 1$.

Warning: If you have more than one character in the exponent, you must use curly braces for grouping. Correct: x^{2n} . Wrong: x^2n .

Subscripts

Index: x_i and $2x_i + 1$.

Beware: curly braces needed for grouping, as before.

Correct: x_{2j+1} . Wrong: x_{2j+1} .

Index: x_i and $2x_i + 1$.

Beware: curly braces needed for grouping, as before. Correct: x_{2j+1} . Wrong: $x_{2j} + 1$.

Equations

$1+1 = 2$. $x \neq y$.

$5 < 6$. $5 \leq 7$. $5 \geq 0$.

$x \in S$. $y \notin S$. $S \subseteq T$.

$1 + 1 = 2$. $x \neq y$.

$5 < 6$. $5 \leq 7$. $5 \geq 0$.

$x \in S$. $y \notin S$. $S \subseteq T$.

Logic

$\neg P$. $P \vee Q$. $R \wedge S$.

$T \implies U$. $P \vee P \equiv P$.

I claim that $\forall x \in S. P(x)$.

Moreover, $\exists x \in S. Q(x)$.

$\neg P. P \vee Q. R \wedge S$.

$T \implies U. P \vee P \equiv P$.

I claim that $\forall x \in S. P(x)$. Moreover, $\exists x \in S. Q(x)$.

Some examples

Consider any integer $n > 2$.
Then the equation $x^n + y^n = z^n$
has no solutions for x, y, z in the
integers.

Consider any integer $n > 2$. Then the equation $x^n + y^n = z^n$ has no solutions for x, y, z in the integers.

Some examples

Define $f(1) = 1$,
 $f(n) = f(2n+1)$ if n is odd,
and $f(n) = f(n/2)$ if n is even.
Goldbach conjectured that $f(n)$
always terminates and returns 1.

Define $f(1) = 1$, $f(n) = f(2n + 1)$ if n is odd, and
 $f(n) = f(n/2)$ if n is even. Goldbach conjectured
that $f(n)$ always terminates and returns 1.

Displayed equations

Sometimes you want an equation on its own line, like this:

```
\[ (x-3)^2 \ge 0. \]
```

Sometimes you want an equation on its own line, like this:

$$(x - 3)^2 \geq 0.$$

Sums, products, and fractions

Useful:

$$\backslash[1^2 + 2^2 + \dots + n^2$$

$$= \sum_{i=1}^n i^2$$

$$= \frac{n(n+1)(2n+1)}{6}. \backslash]$$

$$\backslash[n! = \prod_{j=1}^n j. \backslash]$$

Useful:

$$1^2 + 2^2 + \dots + n^2 = \sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}.$$

$$n! = \prod_{j=1}^n j.$$

Environments

```
\begin{quote}
```

```
Q: How many Stanford students does it take  
to screw in a light bulb?
```

```
A: One, dude.
```

```
\end{quote}
```

Q: How many Stanford students does it take
to screw in a light bulb?

A: One, dude.

Environments

```
\begin{verbatim}
```

```
Q: How many professors does it take to  
screw in a light bulb?
```

```
A: Only one, but they get three tech.  
reports out of it.
```

```
\end{verbatim}
```

```
Q: How many professors does it  
take to screw in a light bulb?
```

```
A: Only one, but they get three  
tech. reports out of it.
```

Environments

```
\begin{center}
```

Q: How many slides does it take until we get to a joke that is actually funny?

A: ∞ .

```
\end{center}
```

Q: How many slides does it take until we get to a joke that is actually funny?

A: ∞ .

Lists

How do you catch a blue elephant?

```
\begin{itemize}
```

```
\item With a blue elephant net, of course.
```

```
\item But it better be a friggin' big net.
```

```
\end{itemize}
```

How do you catch a blue elephant?

- Use a blue elephant net, of course.
- But it better be a friggin' big net.

Numbered lists

How do you catch a red elephant?

```
\begin{enumerate}
```

```
\item Hold his nose until he turns blue.
```

```
\item Then use a blue elephant net.
```

```
\item[(iv)] Hey, my younger brother  
thought it was funny. Once.
```

```
\end{enumerate}
```

How do you catch a red elephant?

1. Hold his nose until he turns blue.

2. Then use a blue elephant net.

(iv) Hey, my younger brother thought it was funny.
Once.

Multi-line equations

```
\begin{align*}
(x-y)(x+y) &= x^2-y^2 \\
&= x^2-9+9-y^2 \\
&= (x-3)(x+3) + (3-y)(3+y)
\end{align*}
```

$$\begin{aligned}
(x-y)(x+y) &= x^2 - y^2 \\
&= x^2 - 9 + 9 - y^2 \\
&= (x-3)(x+3) + (3-y)(3+y)
\end{aligned}$$

Equations with justifications

```
\begin{align*}
f(n) &= f(n-1) + n \tag{by defn of $f$} \\
&= (n-1)n/2 + n \tag{by inductive hyp} \\
&= n(n+1)/2 \tag{simple algebra} \\
\end{align*}
```

$$\begin{aligned} f(n) &= f(n-1) + n && \text{(by defn of } f) \\ &= (n-1)n/2 + n && \text{(by inductive hyp)} \\ &= n(n+1)/2 && \text{(simple algebra)} \end{aligned}$$

Theorems and proofs

```
\begin{theorem} A ham sandwich is better  
than good sex. \end{theorem}  
\begin{proof} A ham sandwich is better  
than nothing. Also, nothing is better  
than good sex. The result follows by  
transitivity. \end{proof}
```

Theorem 1. *A ham sandwich is better than good sex.*

Proof. A ham sandwich is better than nothing. Also, nothing is better than good sex. The result follows by transitivity. \square

Pitfalls

Beware: Some characters are special, and can't be used from within text mode. This includes `\%$#&_{}<>~^.`

Many of them can be produced by prepending a backslash. For instance, `55%` is produced by typing `"55\"%`, and `{1,2,3}` by `"$\{1,2,3\}$"`.

Error messages

```
Since  $x^2=1$ , we know  $x=1$  or  $x=-1$ .
```

```
This is TeX, Version 3.14159 (Web2C 7.3.1)
```

```
...
```

```
! Missing $ inserted.
```

```
<inserted text>
```

```
      $
```

```
1.25 Since  $x^$ 
```

```
       $2=1$ , we know  $x=1$  or  $x=-1$ .
```

```
?
```

Type “h” for help, then “x” to exit. Next, go look at line 25 of the source document. \LaTeX is telling you that there is a “\$” missing somewhere near there.

Summary

L^AT_EX is cool stuff. Give it a try.

We'll have resources posted on the web page.

And ask us if you have questions.