


Regular B-Spline Subdivision



$$\mathbf{P}_{11} = \mathbf{H}_{B1} \mathbf{P} \mathbf{H}_{B1}^T$$


In this parametric view these knot points are collocated.

The 3D control points are not.

17

17

Regular B-Spline Subdivision




$$\mathbf{P}_{11} = \mathbf{H}_{B1} \mathbf{P} \mathbf{H}_{B1}^T$$

$$\mathbf{P}_{12} = \mathbf{H}_{B1} \mathbf{P} \mathbf{H}_{B2}^T$$

18

18

Regular B-Spline Subdivision




$$\mathbf{P}_{11} = \mathbf{H}_{B1} \mathbf{P} \mathbf{H}_{B1}^T$$

$$\mathbf{P}_{12} = \mathbf{H}_{B1} \mathbf{P} \mathbf{H}_{B2}^T$$

$$\mathbf{P}_{22} = \mathbf{H}_{B2} \mathbf{P} \mathbf{H}_{B2}^T$$

Regular B-Spline Subdivision



$$\mathbf{P}_{11} = \mathbf{H}_{B1} \mathbf{P} \mathbf{H}_{B1}^T$$


$$\mathbf{P}_{12} = \mathbf{H}_{B1} \mathbf{P} \mathbf{H}_{B2}^T$$

$$\mathbf{P}_{22} = \mathbf{H}_{B2} \mathbf{P} \mathbf{H}_{B2}^T$$

$$\mathbf{P}_{21} = \mathbf{H}_{B2} \mathbf{P} \mathbf{H}_{B1}^T$$

$$\mathbf{H}_{B1} = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ \frac{1}{8} & \frac{3}{4} & \frac{1}{8} & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & \frac{1}{8} & \frac{3}{4} & \frac{1}{8} \end{bmatrix} \quad \mathbf{H}_{B2} = \begin{bmatrix} \frac{1}{8} & \frac{3}{4} & \frac{1}{8} & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & \frac{1}{8} & \frac{3}{4} & \frac{1}{8} \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix}$$

Regular B-Spline Subdivision



$$\mathbf{P}^{i+1} = \mathbf{H} \mathbf{P}^i$$


Length 25 vector of fine CPs

25 x 16 subdivision matrix

Length 16 vector of coarse CP

21

Regular B-Spline Subdivision



$$\mathbf{P}^{i+1} = \mathbf{H} \mathbf{P}^i$$


Inspection would reveal a pattern

- Face points
- Edge points
- Vertex points

22


Regular B-Spline Subdivision

Face point




$$f = \frac{v_1 + v_2 + v_3 + v_4}{4}$$

Edge point



$$e = \frac{v_1 + v_2 + f_1 + f_2}{4}$$

Vertex point




$$v = \frac{f_1 + f_2 + f_3 + f_4 + 2(m_1 + m_2 + m_3 + m_4) + 4p}{16}$$

m midpoint of edge, not "edge point"
 p old "vertex point"

23

Regular B-Spline Subdivision


- Recall that control mesh approaches surface



24

Regular B-Spline Subdivision

- Limit of subdivision is the surface




Irregular B-Spline Subdivision

- Catmull-Clark Subdivision
- Generalizes regular B-Spline subdivision

An irregular patch

Non-quad face

Extraordinary vertex



Irregular B-Spline Subdivision

- Catmull-Clark Subdivision
 - Generalizes regular B-Spline subdivision
 - Rules reduce to regular for ordinary vertices/faces

f = average of surrounding vertices

$$e = \frac{f_1 + f_2 + v_1 + v_2}{4}$$

$$v = \frac{\bar{f}}{n} + \frac{2\bar{m}}{n} + \frac{p(n-3)}{n}$$


\bar{m} = average of adjacent midpoints

\bar{f} = average of adjacent face points

n = valence of vertex


27

Catmull-Clark Subdivision




28

Catmull-Clark Subdivision




29

Catmull-Clark Subdivision



30

Catmull-Clark Subdivision



31

31

Continuity of Catmull-Clark

- In “ordinary” regions
 - Surface is fully C^2 everywhere except extraordinary points
 - Fast evaluation by matrix exponentiation
 - See “Exact Evaluation Of Catmull-Clark Subdivision Surfaces At Arbitrary Parameter Values” by Jos Stam, SIGGRAPH 1998.
- At extraordinary points
 - Surface is at least C^1
 - Curvature is Lipschitz continuous at extraordinary points

32

32
