

# CS-184: Computer Graphics

## Lecture #5: 3D Transformations and Rotations

Prof. James O'Brien  
University of California, Berkeley

1/20/14 09:10

---

---

---

---

---

---

---

---

---

---

---

---

### Today

- Transformations in 3D
- Rotations
  - Matrices
  - Euler angles
  - Exponential maps
  - Quaternions

2

---

---

---

---

---

---

---

---

---

---

---

---

## 3D Transformations

- Generally, the extension from 2D to 3D is straightforward
  - Vectors get longer by one
  - Matrices get extra column and row
  - SVD still works the same way
  - Scale, Translation, and Shear all basically the same
- Rotations get interesting

3

## Translations

$$\tilde{\mathbf{A}} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \quad \text{For 2D}$$

$$\tilde{\mathbf{A}} = \begin{bmatrix} 1 & 0 & 0 & t_x \\ 0 & 1 & 0 & t_y \\ 0 & 0 & 1 & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad \text{For 3D}$$

4

Scales	
$\tilde{\mathbf{A}} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix}$	For 2D
$\tilde{\mathbf{A}} = \begin{bmatrix} s_x & 0 & 0 & 0 \\ 0 & s_y & 0 & 0 \\ 0 & 0 & s_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$	For 3D (Axis-aligned!)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

Shears	
$\tilde{\mathbf{A}} = \begin{bmatrix} 1 & h_{xy} & 0 \\ h_{yx} & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	For 2D
$\tilde{\mathbf{A}} = \begin{bmatrix} 1 & h_{xy} & h_{xz} & 0 \\ h_{yx} & 1 & h_{yz} & 0 \\ h_{zx} & h_{zy} & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$	For 3D (Axis-aligned!)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


## Shears

$$\tilde{\mathbf{A}} = \begin{bmatrix} 1 & h_{xy} & h_{xz} & 0 \\ h_{yx} & 1 & h_{yz} & 0 \\ h_{zx} & h_{zy} & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Shears y into x


## Rotations

- 3D Rotations fundamentally more complex than in 2D
- 2D: amount of rotation
- 3D: amount and axis of rotation



## Rotations


- Rotations still orthonormal
- $\text{Det}(\mathbf{R}) = 1 \neq -1$
- Preserve lengths and distance to origin
- 3D rotations DO NOT COMMUTE!
- Right-hand rule **DO NOT COMMUTE!**
- Unique matrices



9

## Axis-aligned 3D Rotations

- 2D rotations implicitly rotate about a third out of plane axis



10

## Axis-aligned 3D Rotations

- 2D rotations implicitly rotate about a third out of plane axis

$$\mathbf{R} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} \quad \mathbf{R} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



Note: looks same as  $\tilde{\mathbf{R}}$



11


## Axis-aligned 3D Rotations

$$\mathbf{R}_z = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{bmatrix}$$

$$\mathbf{R}_y = \begin{bmatrix} \cos(\theta) & 0 & \sin(\theta) \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{bmatrix}$$

$$\mathbf{R}_x = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

"Z is in your face"




12

## Axis-aligned 3D Rotations

$$\mathbf{R}_x = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{bmatrix}$$

Also right handed "Zup"

$$\mathbf{R}_y = \begin{bmatrix} \cos(\theta) & 0 & \sin(\theta) \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{bmatrix}$$



$$\mathbf{R}_z = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

13

## Axis-aligned 3D Rotations

• Also known as "direction-cosine" matrices

$$\mathbf{R}_x = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{bmatrix} \quad \mathbf{R}_y = \begin{bmatrix} \cos(\theta) & 0 & \sin(\theta) \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{bmatrix}$$

$$\mathbf{R}_z = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

14

## Arbitrary Rotations


- Can be built from axis-aligned matrices:

$$\mathbf{R} = \mathbf{R}_{\hat{z}} \cdot \mathbf{R}_{\hat{y}} \cdot \mathbf{R}_{\hat{x}}$$

- Result due to Euler... hence called Euler Angles
- Easy to store in vector

- But NOT a vector:


$$\mathbf{R} = \text{rot}(x, y, z)$$



15

## Arbitrary Rotations

$$\mathbf{R} = \mathbf{R}_{\hat{z}} \cdot \mathbf{R}_{\hat{y}} \cdot \mathbf{R}_{\hat{x}}$$




16



## Arbitrary Rotations

- Allows tumbling
- Euler angles are non-unique
- Gimbal-lock
- Moving -vs- fixed axes
  - Reverse of each other




17

## Exponential Maps

- Direct representation of arbitrary rotation
- AKA: axis-angle, angular displacement vector
- Rotate  $\theta$  degrees about some axis
- Encode  $\theta$  by length of vector

$$\theta = |\mathbf{r}|$$




18

## Exponential Maps

- Given vector  $\mathbf{r}$ , how to get matrix  $\mathbf{R}$
- Method from text:
  1. rotate about  $x$  axis to put  $\mathbf{r}$  into the  $x$ - $y$  plane
  2. rotate about  $z$  axis align  $\mathbf{r}$  with the  $x$  axis
  3. rotate  $\theta$  degrees about  $x$  axis
  4. undo #2 and then #1
  5. composite together

19


## Exponential Maps



- Vector expressing a point has two parts
  - $\mathbf{x}_{\parallel}$  does not change
  - $\mathbf{x}_{\perp}$  rotates like a 2D point

20


## Exponential Maps



## Exponential Maps

• Rodriguez Formula

$$\begin{aligned}
 \mathbf{x}' = & \hat{\mathbf{r}}(\hat{\mathbf{r}} \cdot \mathbf{x}) \\
 & + \sin(\theta)(\hat{\mathbf{r}} \times \mathbf{x}) \\
 & - \cos(\theta)(\hat{\mathbf{r}} \times (\hat{\mathbf{r}} \times \mathbf{x}))
 \end{aligned}$$



Linear in  $\mathbf{x}$

Actually a minor variation ... 22

## Exponential Maps

- Building the matrix

$$\mathbf{x}' = ((\hat{\mathbf{r}}\hat{\mathbf{r}}^t) + \sin(\theta)(\hat{\mathbf{r}}\times) - \cos(\theta)(\hat{\mathbf{r}}\times)(\hat{\mathbf{r}}\times))\mathbf{x}$$

$$(\hat{\mathbf{r}}\times) = \begin{bmatrix} 0 & -\hat{r}_z & \hat{r}_y \\ \hat{r}_z & 0 & -\hat{r}_x \\ -\hat{r}_y & \hat{r}_x & 0 \end{bmatrix}$$

Antisymmetric matrix

$$(\mathbf{a}\times)\mathbf{b} = \mathbf{a}\times\mathbf{b}$$

Easy to verify by expansion

23


## Exponential Maps

- Allows tumbling
- No gimbal-lock!
- Orientations are space within  $\pi$ -radius ball
- Nearly unique representation
- Singularities on shells at  $2\pi$
- Nice for interpolation

24

## Exponential Maps

• Why exponential?



• Instead of rotating once by  $\theta$ , let's do  $n$  small rotations of  $\theta/n$

• Now the angle is small, so the rotated  $\mathbf{x}$  is approximately

$$\mathbf{x} + (\theta/n)\hat{\mathbf{r}} \times \mathbf{x} \\ = \left( \mathbf{I} + \frac{(\hat{\mathbf{r}} \times)\theta}{n} \right) \mathbf{x}$$

• Do it  $n$  times and you get

$$\mathbf{x}' = \left( \mathbf{I} + \frac{(\hat{\mathbf{r}} \times)\theta}{n} \right)^n \mathbf{x}$$

## Exponential Maps

$$\mathbf{x}' = \lim_{n \rightarrow \infty} \left( \mathbf{I} + \frac{(\hat{\mathbf{r}} \times)\theta}{n} \right)^n \mathbf{x}$$

• Remind you of anything?

$$\lim_{n \rightarrow \infty} \left( 1 + \frac{a}{n} \right)^n \text{ is a definition of } e^a$$

• So the rotation we want is the exponential of  $(\hat{\mathbf{r}} \times)\theta$ !

• In fact you can just plug it into the infinite series...

## Exponential Maps

- Why exponential?
- Recall series expansion of  $e^x$

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

27

## Exponential Maps

- Why exponential?
- Recall series expansion of  $e^x$
- Euler: what happens if you put in  $i\theta$  for  $x$

$$\begin{aligned} e^{i\theta} &= 1 + \frac{i\theta}{1!} + \frac{-\theta^2}{2!} + \frac{-i\theta^3}{3!} + \frac{\theta^4}{4!} + \dots \\ &= \left(1 + \frac{-\theta^2}{2!} + \frac{\theta^4}{4!} + \dots\right) + i\left(\frac{\theta}{1!} + \frac{-\theta^3}{3!} + \dots\right) \\ &= \cos(\theta) + i\sin(\theta) \end{aligned}$$

28

## Exponential Maps

• Why exponential?

$$e^{(\hat{\mathbf{r}} \times) \theta} = \mathbf{I} + \frac{(\hat{\mathbf{r}} \times) \theta}{1!} + \frac{(\hat{\mathbf{r}} \times)^2 \theta^2}{2!} + \frac{(\hat{\mathbf{r}} \times)^3 \theta^3}{3!} + \frac{(\hat{\mathbf{r}} \times)^4 \theta^4}{4!} + \dots$$

But notice that:  $(\hat{\mathbf{r}} \times)^3 = -(\hat{\mathbf{r}} \times)$

$$e^{(\hat{\mathbf{r}} \times) \theta} = \mathbf{I} + \frac{(\hat{\mathbf{r}} \times) \theta}{1!} + \frac{(\hat{\mathbf{r}} \times)^2 \theta^2}{2!} - \frac{(\hat{\mathbf{r}} \times) \theta^3}{3!} + \frac{(\hat{\mathbf{r}} \times)^2 \theta^4}{4!} + \dots$$

29

## Exponential Maps

$$e^{(\hat{\mathbf{r}} \times) \theta} = \mathbf{I} + \frac{(\hat{\mathbf{r}} \times) \theta}{1!} + \frac{(\hat{\mathbf{r}} \times)^2 \theta^2}{2!} - \frac{(\hat{\mathbf{r}} \times) \theta^3}{3!} + \frac{(\hat{\mathbf{r}} \times)^2 \theta^4}{4!} + \dots$$

$$e^{(\hat{\mathbf{r}} \times) \theta} = (\hat{\mathbf{r}} \times) \left( \frac{\theta}{1!} - \frac{\theta^3}{3!} + \dots \right) + \mathbf{I} + (\hat{\mathbf{r}} \times)^2 \left( \frac{\theta^2}{2!} - \frac{\theta^4}{4!} + \dots \right)$$

$$e^{(\hat{\mathbf{r}} \times) \theta} = (\hat{\mathbf{r}} \times) \sin(\theta) + \mathbf{I} + (\hat{\mathbf{r}} \times)^2 (1 - \cos(\theta))$$

30

## Quaternions

- More popular than exponential maps
- Natural extension of  $e^{i\theta} = \cos(\theta) + i \sin(\theta)$
- Due to Hamilton (1843)
  - Interesting history
  - Involves "hermaphroditic monsters"

31

## Quaternions

- Uber-Complex Numbers

$$q = (z_1, z_2, z_3, s) = (\mathbf{z}, s)$$

$$q = iz_1 + jz_2 + kz_3 + s$$

$$i^2 = j^2 = k^2 = -1 \quad \begin{array}{ll} ij = k & ji = -k \\ jk = i & kj = -i \\ ki = j & ik = -j \end{array}$$

32



## Quaternions

- Multiplication natural consequence of defn.

$$\mathbf{q} \cdot \mathbf{p} = (\mathbf{z}_q s_p + \mathbf{z}_p s_q + \mathbf{z}_p \times \mathbf{z}_q, s_p s_q - \mathbf{z}_p \cdot \mathbf{z}_q)$$

- Conjugate

$$\mathbf{q}^* = (-\mathbf{z}, s)$$

- Magnitude

$$\|\mathbf{q}\|^2 = \mathbf{z} \cdot \mathbf{z} + s^2 = \mathbf{q} \cdot \mathbf{q}^*$$

33

## Quaternions

- Vectors as quaternions

$$\mathbf{v} = (\mathbf{v}, 0)$$

- Rotations as quaternions

$$\mathbf{r} = (\hat{\mathbf{r}} \sin \frac{\theta}{2}, \cos \frac{\theta}{2})$$

- Rotating a vector

$$\mathbf{x}' = \mathbf{r} \cdot \mathbf{x} \cdot \mathbf{r}^*$$

- Composing rotations

$$\mathbf{r} = \mathbf{r}_1 \cdot \mathbf{r}_2 \quad \text{Compare to Exp. Map}$$

34

# Quaternions

- No tumbling
- No gimbal-lock
- Orientations are “double unique”
- Surface of a 3-sphere in 4D  $||r|| = 1$
- Nice for interpolation

28

---

---

---

---

---

---

---

---


---

---

---

---

# Interpolation



29

---

---

---

---

---

---

---

---

---

---

---

---

## Rotation Matrices

- Eigen system
  - One real eigenvalue
  - Real axis is axis of rotation
  - Imaginary values are 2D rotation as complex number
- Logarithmic formula

$$(\hat{\mathbf{r}} \times) = \ln(\mathbf{R}) = \frac{\theta}{2 \sin \theta} (\mathbf{R} - \mathbf{R}^T)$$

$$\theta = \cos^{-1} \left( \frac{\text{Tr}(\mathbf{R}) - 1}{2} \right)$$

Similar formulae as for exponential... 27


## Rotation Matrices

- Consider:

$$\mathbf{R}\mathbf{I} = \begin{bmatrix} r_{xx} & r_{xy} & r_{xz} \\ r_{yx} & r_{yy} & r_{yz} \\ r_{zx} & r_{zy} & r_{zz} \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Columns are coordinate axes after (true for general matrices)
- Rows are original axes in original system (not true for general matrices)

28




---

---

---

---

---

---

---

---

## Scene Graphs

- Draw scene with pre-and-post-order traversal
  - Apply node, draw children, undo node if applicable
- Nodes can do pretty much anything
  - Geometry, transformations, groups, color, switch, scripts, etc.
  - Node types are application/implementation specific
- Requires a stack to implement "undo" post children
- Nodes can cache their children
- Instances make it a DAG, not strictly a tree
- Will use these trees later for bounding box trees

---

---

---


---

---

---

---

---




---



---



---



---



---



---



---



---