



CS-184: Computer Graphics Lecture #21: Integration Basics Prof. James O'Brien University of California, Berkeley Venerated	1	
Today	2	
Introduction to Simulation Basic particle systems Time integration (simple version)		
	1	









5 Particle Systems • Single particles are very simple Large groups can produce interesting effects Supplement basic ballistic rules Collisions Interactions Force fields Springs Others... 6 PARTICLE DREAMS Karl Sims Optomystic

Particle Systems

- · Single particles are very simple
- · Large groups can produce interesting effects
- Supplement basic ballistic rules
- Collisions
- Interactions
- Force fields
- Springs
- Others...



Basic Particles

- Basic governing equation
- ${}^{\star}oldsymbol{f}$ is a sum of a number of things

- $m{f}$ is a sum of a number of things $\ddot{m{x}} = rac{1}{m} m{f}$ Gravity: constant downward force proportional to mass
- Simple drag: force proportional to negative velocity
- Particle interactions: particles mutually attract and/or repell
- * Beware $O(n^2)$ complexity!
- Force fields
- Wind forces
- User interaction

9 Basic Particles Properties other than position • Color Temp Age • Differential equations also needed to govern these properties Collisions and other constrains directly modify position and/or velocity 10 Particle Rules Multiple Burst Bryan E. Feldman, James F. O'Brien, and Okan Arikan. "Animating Suspended Particle Explosions". In Proceedings of ACM S/GGRAPH 2003, pages 708–715, August 2003.

- Euler's Method
- Simple
- Commonly used
- Very inaccurate
- Most often goes unstable

$$\mathbf{x}^{t+\Delta t} = \mathbf{x}^t + \Delta t \, \mathbf{\dot{x}}^t$$

$$\mathbf{\dot{x}}^{t+\Delta t} = \mathbf{\dot{x}}^t + \Delta t \, \mathbf{\ddot{x}}^t$$

...

11

Integration

• For now let's pretend

$$\boldsymbol{f} = m\boldsymbol{v}$$

· Velocity (rather than acceleration) is a function of force



 $\dot{\boldsymbol{x}} = f(\boldsymbol{x}, t)$


Witkin and Baraff

Note: Second order ODEs can be turned into first order ODEs using extra variables.

• For now let's pretend

$$\boldsymbol{f} = m\boldsymbol{v}$$

• Velocity (rather than acceleration) is a function of force




$$\dot{\boldsymbol{x}} = \mathsf{f}(\boldsymbol{x},t)$$

13

14

Integration

- With numerical integration, errors accumulate
- Euler integration is particularly bad



$$x := x + \Delta t \ \mathsf{f}(\boldsymbol{x}, t)$$

(, ,

- Stability issues can also arise
- Occurs when errors lead to larger errors
- Often more serious than error issues





 $\dot{\boldsymbol{x}} = [-\sin(\omega t) , -\cos(\omega t)]$

15

Integration


Modified Euler

$$\boldsymbol{x}^{t+\Delta t} = \boldsymbol{x}^t + \frac{\Delta t}{2} \left(\dot{\boldsymbol{x}}^t + \dot{\boldsymbol{x}}^{t+\Delta t} \right)$$

$$\dot{\boldsymbol{x}}^{t+\Delta t} = \dot{\boldsymbol{x}}^t + \Delta t \ \ddot{\boldsymbol{x}}^t$$

$$oldsymbol{x}^{t+\Delta t} = oldsymbol{x}^t + \Delta t \ \dot{oldsymbol{x}}^t + rac{(\Delta t)^2}{2} \ \ddot{oldsymbol{x}}^t$$

- Midpoint method
- a. Compute half Euler step
- b. Eval. derivative at halfway
- c. Retake step
- Other methods
- Verlet
- Runge-Kutta
- · And many others...



17

Integration

- Implicit methods
- Informally (incorrectly) called backward methods
- Use derivatives in the future for the current step

$$\begin{split} \boldsymbol{x}^{t+\Delta t} &= \boldsymbol{x}^t + \Delta t \ \dot{\boldsymbol{x}}^{t+\Delta t} \\ \dot{\boldsymbol{x}}^{t+\Delta t} &= \dot{\boldsymbol{x}}^t + \Delta t \ \ddot{\boldsymbol{x}}^{t+\Delta t} \\ \\ \dot{\boldsymbol{x}}^{t+\Delta t} &= \mathsf{V}(\boldsymbol{x}^{t+\Delta t}, \dot{\boldsymbol{x}}^{t+\Delta t}, t + \Delta t) \\ \\ \ddot{\boldsymbol{x}}^{t+\Delta t} &= \mathsf{A}(\boldsymbol{x}^{t+\Delta t}, \dot{\boldsymbol{x}}^{t+\Delta t}, t + \Delta t) \end{split}$$

- Implicit methods
- · Informally (incorrectly) called backward methods
- Use derivatives in the future for the current step

$$\begin{split} \dot{\boldsymbol{x}}^{t+\Delta t} &= \dot{\boldsymbol{x}}^t + \Delta t \; \mathsf{V}(\boldsymbol{x}^{t+\Delta t}, \dot{\boldsymbol{x}}^{t+\Delta t}, t + \Delta t) \\ \dot{\boldsymbol{x}}^{t+\Delta t} &= \dot{\boldsymbol{x}}^t + \Delta t \; \mathsf{A}(\boldsymbol{x}^{t+\Delta t}, \dot{\boldsymbol{x}}^{t+\Delta t}, t + \Delta t) \end{split}$$

- * Solve nonlinear problem for $~m{x}^{t+\Delta t}$ and $\dot{m{x}}^{t+\Delta t}$
- · This is fully implicit backward Euler
- Many other implicit methods exist...
- Modified Euler is *partially* implicit as is Verlet

19

Temp Slide





Need to draw reverse diagrams....

- Semi-Implicit
- · Approximate with linearized equations

$$\mathsf{V}(\boldsymbol{x}^{t+\Delta t}, \dot{\boldsymbol{x}}^{t+\Delta t}) \approx \mathsf{V}(\boldsymbol{x}^t, \dot{\boldsymbol{x}}^t) + \mathbf{A} \cdot (\Delta \boldsymbol{x}) + \mathbf{B} \cdot (\Delta \dot{\boldsymbol{x}})$$

$$\mathsf{A}(\boldsymbol{x}^{t+\Delta t}, \dot{\boldsymbol{x}}^{t+\Delta t}) \approx \mathsf{A}(\boldsymbol{x}^t, \dot{\boldsymbol{x}}^t) + \mathbf{C} \cdot (\Delta \boldsymbol{x}) + \mathbf{D} \cdot (\Delta \dot{\boldsymbol{x}})$$


$$\begin{bmatrix} \boldsymbol{x}^{t+\Delta t} \\ \dot{\boldsymbol{x}}^{t+\Delta t} \end{bmatrix} = \begin{bmatrix} \boldsymbol{x}^t \\ \dot{\boldsymbol{x}}^t \end{bmatrix} + \Delta t \left(\begin{bmatrix} \dot{\boldsymbol{x}}^t \\ \ddot{\boldsymbol{x}}^t \end{bmatrix} + \begin{bmatrix} \mathbf{A} \ \mathbf{B} \\ \mathbf{C} \ \mathbf{D} \end{bmatrix} \begin{bmatrix} \Delta \boldsymbol{x} \\ \Delta \dot{\boldsymbol{x}} \end{bmatrix} \right)$$

21

Integration

- Explicit methods can be conditionally stable
- Depends on time-step and **stiffness** of system
- Fully implicit can be **un**conditionally stable
- May still have large errors
- Semi-implicit can be conditionally stable
- Nonlinearities can cause instability
- Generally more stable than explicit
- · Comparable errors as explicit
- · Often show up as excessive damping

- Integrators can be analyzed in modal domain
- System have different component behaviors
- Integrators impact components differently



23

23

Suggested Reading

- Physically Based Modeling: Principles and Practice
- · Andy Witkin and David Baraff
- http://www-2.cs.cmu.edu/~baraff/sigcourse/index.html
- Numerical Recipes in C++
- Chapter 16
- Any good text on integrating ODE's