


# BAYESIAN NETWORKS

## CHAPTER 14.1-3

### Example

Topology of network encodes conditional independence assertions:



*Weather* is independent of the other variables

*Toothache* and *Catch* are conditionally independent given *Cavity*

### Outline

- ◇ Syntax
- ◇ Semantics
- ◇ Parameterized distributions

### Example

I'm at work, neighbor John calls to say my alarm is ringing, but neighbor Mary doesn't call. Sometimes it's set off by minor earthquakes. Is there a burglar?

Variables: *Burglar*, *Earthquake*, *Alarm*, *JohnCalls*, *MaryCalls*

Network topology reflects "causal" knowledge:

- A burglar can set the alarm off
- An earthquake can set the alarm off
- The alarm can cause Mary to call
- The alarm can cause John to call

### Bayesian networks


A simple, graphical notation for conditional independence assertions and hence for compact specification of full joint distributions

Syntax:

- a set of nodes, one per variable
- a directed, acyclic graph (link  $\approx$  "directly influences")
- a conditional distribution for each node given its parents:  
 $P(X_i | Parents(X_i))$

In the simplest case, conditional distribution represented as a conditional probability table (CPT) giving the distribution over  $X_i$  for each combination of parent values

### Example contd.



## Compactness


A CPT for Boolean  $X_i$  with  $k$  Boolean parents has  $2^k$  rows for the combinations of parent values

Each row requires one number  $p$  for  $X_i = \text{true}$  (the number for  $X_i = \text{false}$  is just  $1 - p$ )

If each variable has no more than  $k$  parents, the complete network requires  $O(n \cdot 2^k)$  numbers

I.e., grows linearly with  $n$ , vs.  $O(2^n)$  for the full joint distribution


For burglary net,  $1 + 1 + 4 + 2 + 2 = 10$  numbers (vs.  $2^5 - 1 = 31$ )



Chapter 14.1-3 7

## Local semantics

Local semantics: each node is conditionally independent of its nondescendants given its parents



Theorem: Local semantics  $\Leftrightarrow$  global semantics

Chapter 14.1-3 10


## Global semantics

Global semantics defines the full joint distribution as the product of the local conditional distributions:

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

e.g.,  $P(j \wedge m \wedge a \wedge \neg b \wedge \neg e)$

=




Chapter 14.1-3 8

## Markov blanket

Each node is conditionally independent of all others given its

Markov blanket: parents + children + children's parents



Chapter 14.1-3 11


## Global semantics

"Global" semantics defines the full joint distribution as the product of the local conditional distributions:

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

e.g.,  $P(j \wedge m \wedge a \wedge \neg b \wedge \neg e)$

$$\begin{aligned} &= P(j|a)P(m|a)P(a|\neg b, \neg e)P(\neg b)P(\neg e) \\ &= 0.9 \times 0.7 \times 0.001 \times 0.999 \times 0.998 \\ &\approx 0.00063 \end{aligned}$$



Chapter 14.1-3 9

## Constructing Bayesian networks

Need a method such that a series of locally testable assertions of conditional independence guarantees the required global semantics

1. Choose an ordering of variables  $X_1, \dots, X_n$
2. For  $i = 1$  to  $n$   
add  $X_i$  to the network  
select parents from  $X_1, \dots, X_{i-1}$  such that  
 $P(X_i | \text{Parents}(X_i)) = P(X_i | X_1, \dots, X_{i-1})$   
i.e.,  $X_i$  is conditionally independent of other variables given parents

This choice of parents guarantees the global semantics:

$$\begin{aligned} P(X_1, \dots, X_n) &= \prod_{i=1}^n P(X_i | X_1, \dots, X_{i-1}) \quad (\text{chain rule}) \\ &= \prod_{i=1}^n P(X_i | \text{Parents}(X_i)) \quad (\text{by construction}) \end{aligned}$$

Chapter 14.1-3 12

### Example


Suppose we choose the ordering  $M, J, A, B, E$



$P(J|M) = P(J)$ ?

### Example


Suppose we choose the ordering  $M, J, A, B, E$



$P(J|M) = P(J)$ ? No  
 $P(A|J, M) = P(A|J)$ ?  $P(A|J, M) = P(A)$ ? No  
 $P(B|A, J, M) = P(B|A)$ ? Yes  
 $P(B|A, J, M) = P(B)$ ? No  
 $P(E|B, A, J, M) = P(E|A)$ ?  
 $P(E|B, A, J, M) = P(E|A, B)$ ?

### Example


Suppose we choose the ordering  $M, J, A, B, E$



$P(J|M) = P(J)$ ? No  
 $P(A|J, M) = P(A|J)$ ?  $P(A|J, M) = P(A)$ ?

### Example


Suppose we choose the ordering  $M, J, A, B, E$



$P(J|M) = P(J)$ ? No  
 $P(A|J, M) = P(A|J)$ ?  $P(A|J, M) = P(A)$ ? No  
 $P(B|A, J, M) = P(B|A)$ ? Yes  
 $P(B|A, J, M) = P(B)$ ? No  
 $P(E|B, A, J, M) = P(E|A)$ ? No  
 $P(E|B, A, J, M) = P(E|A, B)$ ? Yes


### Example

Suppose we choose the ordering  $M, J, A, B, E$



$P(J|M) = P(J)$ ? No  
 $P(A|J, M) = P(A|J)$ ?  $P(A|J, M) = P(A)$ ? No  
 $P(B|A, J, M) = P(B|A)$ ?  
 $P(B|A, J, M) = P(B)$ ?


### Example contd.



Deciding conditional independence is hard in noncausal directions  
 (Causal models and conditional independence seem hardwired for humans!)  
 Assessing conditional probabilities is hard in noncausal directions  
 Network is less compact:  $1 + 2 + 4 + 2 + 4 = 13$  numbers needed

### Example: Car diagnosis

Initial evidence: car won't start  
 Testable variables (green), "broken, so fix it" variables (orange)  
 Hidden variables (gray) ensure sparse structure, reduce parameters



### Compact conditional distributions contd.


Noisy-OR distributions model multiple noninteracting causes

- Parents  $U_1 \dots U_k$  include all causes (can add leak node)
- Independent failure probability  $q_i$  for each cause alone  
 $\Rightarrow P(X|U_1 \dots U_j, \neg U_{j+1} \dots \neg U_k) = 1 - \prod_{i=1}^j q_i$

Cold	Flu	Malaria	$P(\text{Fever})$	$P(\neg \text{Fever})$
F	F	F	<b>0.0</b>	1.0
F	F	T	0.9	<b>0.1</b>
F	T	F	0.8	<b>0.2</b>
F	T	T	0.98	$0.02 = 0.2 \times 0.1$
T	F	F	0.4	<b>0.6</b>
T	F	T	0.94	$0.06 = 0.6 \times 0.1$
T	T	F	0.88	$0.12 = 0.6 \times 0.2$
T	T	T	0.988	$0.012 = 0.6 \times 0.2 \times 0.1$


Number of parameters **linear** in number of parents

### Example: Car insurance



### Hybrid (discrete+continuous) networks

Discrete (*Subsidy?* and *Buys?*); continuous (*Harvest* and *Cost*)



- Option 1: discretization—possibly large errors, large CPTs
- Option 2: finitely parameterized canonical families

- Continuous variable, discrete+continuous parents (e.g., *Cost*)
- Discrete variable, continuous parents (e.g., *Buys?*)

### Compact conditional distributions

CPT grows exponentially with number of parents  
 CPT becomes infinite with continuous-valued parent or child

Solution: canonical distributions that are defined compactly

Deterministic nodes are the simplest case:

$$X = f(\text{Parents}(X)) \text{ for some function } f$$

E.g., Boolean functions

$$\text{NorthAmerican} \Leftrightarrow \text{Canadian} \vee \text{US} \vee \text{Mexican}$$

E.g., numerical relationships among continuous variables

$$\frac{\partial \text{Level}}{\partial t} = \text{inflow} + \text{precipitation} - \text{outflow} - \text{evaporation}$$

### Continuous child variables

Need one conditional density function for child variable given continuous parents, for each possible assignment to discrete parents


Most common is the linear Gaussian model, e.g.,:

$$\begin{aligned} P(\text{Cost} = c | \text{Harvest} = h, \text{Subsidy?} = \text{true}) &= N(a_t h + b_t, \sigma_t)(c) \\ &= \frac{1}{\sigma_t \sqrt{2\pi}} \exp\left(-\frac{1}{2} \left(\frac{c - (a_t h + b_t)}{\sigma_t}\right)^2\right) \end{aligned}$$

Mean *Cost* varies linearly with *Harvest*, variance is fixed

Linear variation is unreasonable over the full range  
 but works OK if the **likely** range of *Harvest* is narrow

## Continuous child variables



All-continuous network with LG distributions  
 ⇒ full joint distribution is a multivariate Gaussian


Discrete+continuous LG network is a **conditional Gaussian** network i.e., a multivariate Gaussian over all continuous variables for each combination of discrete variable values

## Discrete variable contd.

Sigmoid (or logit) distribution also used in neural networks:


$$P(\text{Buys?} = \text{true} \mid \text{Cost} = c) = \frac{1}{1 + \exp(-2\frac{-c+\mu}{\sigma})}$$

Sigmoid has similar shape to probit but much longer tails:



## Discrete variable w/ continuous parents

Probability of *Buys?* given *Cost* should be a "soft" threshold:




Probit distribution uses integral of Gaussian:

$$\Phi(x) = \int_{-\infty}^x N(0, 1)(x) dx$$

$$P(\text{Buys?} = \text{true} \mid \text{Cost} = c) = \Phi((-c + \mu)/\sigma)$$

## Why the probit?

1. It's sort of the right shape
2. Can view as hard threshold whose location is subject to noise



## Summary

Bayes nets provide a natural representation for (causally induced) conditional independence

Topology + CPTs = compact representation of joint distribution

Generally easy for (non)experts to construct

Canonical distributions (e.g., noisy-OR) = compact representation of CPTs

Continuous variables ⇒ parameterized distributions (e.g., linear Gaussian)