

## Putnam Mathematical Competition, 2 December 2000

### Problem A1

Let  $A$  be a positive real number. What are the possible values of  $\sum_{j=0}^{\infty} x_j^2$ , given that  $x_0, x_1, x_2, \dots$  are positive numbers for which  $\sum_{j=0}^{\infty} x_j = A$ ?

### Problem A2

Prove that there exist infinitely many integers  $n$  such that  $n$ ,  $n + 1$ , and  $n + 2$  are each the sum of two squares of integers.

[*Example:*  $0 = 0^2 + 0^2$ ,  $1 = 0^2 + 1^2$ , and  $2 = 1^2 + 1^2$ .]

### Problem A3

The octagon  $P_1P_2P_3P_4P_5P_6P_7P_8$  is inscribed in a circle, with the vertices around the circumference in the given order. Given that the polygon  $P_1P_3P_5P_7$  is a square of area 5 and the polygon  $P_2P_4P_6P_8$  is a rectangle of area 4, find the maximum possible area of the octagon.

### Problem A4

Show that the improper integral

$$\lim_{B \rightarrow \infty} \int_0^B \sin(x) \sin(x^2) dx$$

converges.

### Problem A5

Three distinct points with integer coordinates lie in the plane on a circle of radius  $r > 0$ . Show that two of these points are separated by a distance of at least  $r^{1/3}$ .

### Problem A6

Let  $f(x)$  be a polynomial with integer coefficients. Define a sequence  $a_0, a_1, \dots$  of integers such that  $a_0 = 0$  and  $a_{n+1} = f(a_n)$  for all  $n \geq 0$ . Prove that if there exists a positive integer  $m$  for which  $a_m = 0$  then either  $a_1 = 0$  or  $a_2 = 0$ .

**Problem B1**

Let  $a_j, b_j,$  and  $c_j$  be integers for  $1 \leq j \leq N$ . Assume, for each  $j$ , that at least one of  $a_j, b_j, c_j$  is odd. Show that there exist integers  $r, s, t$  such that  $ra_j + sb_j + tc_j$  is odd for at least  $4N/7$  values of  $j, 1 \leq j \leq N$ .

**Problem B2**

Prove that the expression

$$\frac{\gcd(m, n)}{n} \binom{n}{m}$$

is an integer for all pairs of integers  $n \geq m \geq 1$ . [Here  $\binom{n}{m} = \frac{n!}{m!(n-m)!}$  and  $\gcd(m, n)$  is the greatest common divisor of  $m$  and  $n$ .]

**Problem B3**

Let  $f(t) = \sum_{j=1}^N a_j \sin(2\pi jt)$ , where each  $a_j$  is real and  $a_N \neq 0$ . Let  $N_k$  denote the number of zeros (including multiplicities) of  $\frac{d^k f}{dt^k}$ . Prove that

$$N_0 \leq N_1 \leq N_2 \leq \cdots \quad \text{and} \quad \lim_{k \rightarrow \infty} N_k = 2N.$$

**Problem B4**

Let  $f(x)$  be a continuous function such that  $f(2x^2 - 1) = 2xf(x)$  for all  $x$ . Show that  $f(x) = 0$  for  $-1 \leq x \leq 1$ .

**Problem B5**

Let  $S_0$  be a finite set of positive integers. We define finite sets  $S_1, S_2, \dots$  of positive integers as follows:

Integer  $a$  is in  $S_{n+1}$  if and only if exactly one of  $a - 1$  or  $a$  is in  $S_n$ .

Show that there exist infinitely many integers  $N$  for which  $S_N = S_0 \cup \{N + a : a \in S_0\}$ .

**Problem B6**

Let  $B$  be a set of more than  $2^{n+1}/n$  distinct points with coordinates of the form  $(\pm 1, \pm 1, \dots, \pm 1)$  in  $n$ -dimensional space, with  $n \geq 3$ . Show that there are three distinct points in  $B$  which are the vertices of an equilateral triangle.

## Unofficial Solutions

D. J. Bernstein, 3 December 2000

### Problem A1

Let  $A$  be a positive real number. What are the possible values of  $\sum_{j=0}^{\infty} x_j^2$ , given that  $x_0, x_1, x_2, \dots$  are positive numbers for which  $\sum_{j=0}^{\infty} x_j = A$ ?

**Solution:** One can achieve any real number  $s$  with  $0 < s < A^2$  as follows. Define  $u = s/A^2$ ; then  $0 < u < 1$ . Define  $r = (1 - u)/(1 + u)$ ; then  $0 < r < 1$ . Define  $x_j = A(1 - r)r^j$ ; then  $x_j > 0$ . Finally  $\sum x_j = A(1 - r)\sum r^j = A$  and  $\sum x_j^2 = A^2(1 - r)^2 \sum r^{2j} = A^2(1 - r)^2/(1 - r^2) = A^2(1 - r)/(1 + r) = A^2u = s$ .

One cannot achieve any other number, since  $0 < \sum x_j^2 < (\sum x_j)^2 = A^2$ .

### Problem A2

Prove that there exist infinitely many integers  $n$  such that  $n$ ,  $n + 1$ , and  $n + 2$  are each the sum of two squares of integers.

[*Example:*  $0 = 0^2 + 0^2$ ,  $1 = 0^2 + 1^2$ , and  $2 = 1^2 + 1^2$ .]

**Solution:** There are infinitely many integers  $n$  of the form  $2k^2(k + 1)^2$ ; note that  $n = (k^2 + k)^2 + (k^2 + k)^2$ ,  $n + 1 = (k^2 + 2k)^2 + (k^2 - 1)^2$ , and  $n + 2 = (k^2 + k + 1)^2 + (k^2 + k - 1)^2$ .

### Problem A3

The octagon  $P_1P_2P_3P_4P_5P_6P_7P_8$  is inscribed in a circle, with the vertices around the circumference in the given order. Given that the polygon  $P_1P_3P_5P_7$  is a square of area 5 and the polygon  $P_2P_4P_6P_8$  is a rectangle of area 4, find the maximum possible area of the octagon.

**Solution:** The circle circumscribes a square of area 5, so the circle has radius  $\sqrt{5/2}$ . Hence the rectangle has sides  $\sqrt{2}$  and  $\sqrt{8}$ . Without loss of generality assume that  $P_2P_4$  has length  $\sqrt{2}$ .

Put  $P_2, P_4, P_6, P_8$  into the complex plane at  $\sqrt{2}(1/2 + i)$ ,  $\sqrt{2}(-1/2 + i)$ ,  $\sqrt{2}(-1/2 - i)$ ,  $\sqrt{2}(1/2 - i)$ . Put  $P_1$  into the complex plane at  $\sqrt{5/2} \exp(i\theta)$ ; then  $P_3, P_5, P_7$  are at  $i\sqrt{5/2} \exp(i\theta)$ ,  $-\sqrt{5/2} \exp(i\theta)$ ,  $-i\sqrt{5/2} \exp(i\theta)$ .

The triangles  $P_8P_1P_2$  and  $P_4P_5P_6$  each have area  $\sqrt{5} \cos \theta - 1$ . The triangles  $P_2P_3P_4$  and  $P_6P_7P_8$  each have area  $\sqrt{5/4} \cos \theta - 1$ . Hence the octagon has area  $3\sqrt{5} \cos \theta$ . The maximum possible area is  $3\sqrt{5}$ , achieved for  $\theta = 0$ .

### Problem A4

Show that the improper integral

$$\lim_{B \rightarrow \infty} \int_0^B \sin(x) \sin(x^2) dx$$

converges.

**Solution:** Rewrite  $\sin x \sin x^2$  as  $(\cos(x^2 - x) - \cos(x^2 + x))/2$ . In the improper integral  $\int_0^\infty \cos(x^2 + x) dx$  substitute  $u = x^2 + x$  to obtain  $\int_0^\infty 2 \cos u du / (\sqrt{1 + 4u} - 1)$ . The integrand is negative on  $(\pi/2, 3\pi/2)$ , positive on  $(3\pi/2, 5\pi/2)$ , etc. The corresponding integrals form an alternating decreasing series since

$$\int_s^{s+\pi} \frac{2 |\cos u| du}{\sqrt{1 + 4u} - 1} > \int_s^{s+\pi} \frac{2 |\cos u| du}{\sqrt{1 + 4u + 4\pi} - 1} = \int_{s+\pi}^{s+2\pi} \frac{2 |\cos v| dv}{\sqrt{1 + 4v} - 1}.$$

Thus  $\int_0^\infty \cos(x^2 + x) dx$  converges. Similar comments apply to  $\int_0^\infty \cos(x^2 - x) dx$ .

### Problem A5

Three distinct points with integer coordinates lie in the plane on a circle of radius  $r > 0$ . Show that two of these points are separated by a distance of at least  $r^{1/3}$ .

**Solution:** The following solution is stolen from Dave Rusin.

The triangle formed by the points has area  $abc/4r$  where  $a, b, c$  are the distances between the points. If  $a, b, c < r^{1/3}$  then the area is smaller than  $1/4$ ; but the area is at least  $1/2$  since the points have integer coordinates.

### Problem A6

Let  $f(x)$  be a polynomial with integer coefficients. Define a sequence  $a_0, a_1, \dots$  of integers such that  $a_0 = 0$  and  $a_{n+1} = f(a_n)$  for all  $n \geq 0$ . Prove that if there exists a positive integer  $m$  for which  $a_m = 0$  then either  $a_1 = 0$  or  $a_2 = 0$ .

**Solution:** The stated conclusion is false, because the word “either” means that exactly one is true. Presumably the intent was to say that  $a_1 = 0$  or  $a_2 = 0$ .

Fact 1:  $a_{m-1}$  divides  $a_1$ . Proof:  $a_{m-1}$  divides  $f(a_{m-1}) - f(0) = a_m - a_1 = -a_1$ .

Fact 2:  $a_1$  divides  $a_n$  if  $n \geq 0$ . Proof: If  $n = 0$  then  $a_n = 0$ . Otherwise  $a_1$  divides  $a_{n-1}$  by induction, so it divides  $f(a_{n-1}) - f(0) = a_n - a_1$ .

Fact 3:  $a_n - a_{n-1}$  divides  $a_{n+k} - a_{n+k-1}$  if  $n \geq 1$  and  $k \geq 0$ . Proof: If  $k = 0$  then  $a_n - a_{n-1} = a_{n+k} - a_{n+k-1}$ . Otherwise  $a_n - a_{n-1}$  divides  $a_{n+k-1} - a_{n+k-2}$  by induction, so it divides  $f(a_{n+k-1}) - f(a_{n+k-2}) = a_{n+k} - a_{n+k-1}$ .

Fact 4:  $a_n - a_{n-1} \in \{-a_1, a_1\}$  if  $1 \leq n \leq m$ . Proof: Define  $k = m - n$ . Then  $a_n - a_{n-1}$  divides  $a_{n+k} - a_{n+k-1} = a_m - a_{m-1} = -a_{m-1}$ , which divides  $a_1$ ; and  $a_1$  divides  $a_n - a_{n-1}$ .

Fact 5:  $a_2 = 0$ . Proof:  $a_2 - a_1 \in \{-a_1, a_1\}$ . Suppose that  $a_2 \neq 0$ . Then  $a_2 = 2a_1$  and  $a_1 \neq 0$ , so  $m \geq 3$ . Observe that  $a_n = na_1$  for  $n \in \{0, 1, 2\}$ , but not for  $n = m$ . Find the smallest  $n \geq 3$  for which  $a_n \neq na_1$ . Then  $a_{n-1} = (n-1)a_1$ , so  $a_n - a_{n-1} \neq a_1$ , so  $a_n - a_{n-1} = -a_1$ , so  $a_n = (n-2)a_1 = a_{n-2}$ . By induction  $a_k \in \{a_{n-1}, a_{n-2}\}$  for all  $k \geq n$ . In particular  $0 = a_m \in \{a_{n-1}, a_{n-2}\}$ . Thus  $(n-1)a_1 = 0$  or  $(n-2)a_1 = 0$ . Contradiction.

I would have written this problem as follows: “Define  $a_0 = 0$  and  $a_{n+1} = f(a_n)$ , where  $f$  is a polynomial with integer coefficients. Assume that  $a_{2000} = 0$ . Prove that  $a_2 = 0$ .”

### Problem B1

Let  $a_j, b_j,$  and  $c_j$  be integers for  $1 \leq j \leq N$ . Assume, for each  $j$ , that at least one of  $a_j, b_j, c_j$  is odd. Show that there exist integers  $r, s, t$  such that  $ra_j + sb_j + tc_j$  is odd for at least  $4N/7$  values of  $j, 1 \leq j \leq N$ .

**Solution:** Define  $f(u, v, w) = \#\{j : (a_j \bmod 2, b_j \bmod 2, c_j \bmod 2) = (u, v, w)\}$ . Define  $g(r, s, t) = \#\{j : ra_j + sb_j + tc_j \text{ is odd}\}$ . Then

$$\begin{aligned} g(1, 0, 0) &= f(1, 0, 0) + f(1, 0, 1) + f(1, 1, 0) + f(1, 1, 1), \\ g(0, 1, 0) &= f(0, 1, 0) + f(0, 1, 1) + f(1, 1, 0) + f(1, 1, 1), \\ g(0, 0, 1) &= f(0, 0, 1) + f(0, 1, 1) + f(1, 0, 1) + f(1, 1, 1), \\ g(1, 1, 0) &= f(1, 0, 0) + f(0, 1, 0) + f(1, 0, 1) + f(0, 1, 1), \\ g(1, 0, 1) &= f(1, 0, 0) + f(0, 0, 1) + f(1, 1, 0) + f(0, 1, 1), \\ g(0, 1, 1) &= f(0, 1, 0) + f(0, 0, 1) + f(1, 1, 0) + f(1, 0, 1), \\ g(1, 1, 1) &= f(1, 0, 0) + f(0, 1, 0) + f(0, 0, 1) + f(1, 1, 1). \end{aligned}$$

Add:  $g(1, 0, 0) + g(0, 1, 0) + g(0, 0, 1) + g(1, 1, 0) + g(1, 0, 1) + g(0, 1, 1) + g(1, 1, 1) = 4f(1, 0, 0) + 4f(0, 1, 0) + 4f(0, 0, 1) + 4f(1, 1, 0) + 4f(1, 0, 1) + 4f(0, 1, 1) + 4f(1, 1, 1) = 4N$ . Thus  $g(r, s, t) \geq 4N/7$  for some  $(r, s, t)$ .

### Problem B2

Prove that the expression

$$\frac{\gcd(m, n)}{n} \binom{n}{m}$$

is an integer for all pairs of integers  $n \geq m \geq 1$ . [Here  $\binom{n}{m} = \frac{n!}{m!(n-m)!}$  and  $\gcd(m, n)$  is the greatest common divisor of  $m$  and  $n$ .]

**Solution:** Presumably “divisor” means “divisor.”

Find integers  $a, b$  with  $\gcd(m, n) = am + bn$ . Then  $(\gcd(m, n)/n) \binom{n}{m} = a \binom{n-1}{m-1} + b \binom{n}{m}$ .

### Problem B3

Let  $f(t) = \sum_{j=1}^N a_j \sin(2\pi jt)$ , where each  $a_j$  is real and  $a_N \neq 0$ . Let  $N_k$  denote the number of zeros (including multiplicities) of  $\frac{d^k f}{dt^k}$ . Prove that

$$N_0 \leq N_1 \leq N_2 \leq \cdots \quad \text{and} \quad \lim_{k \rightarrow \infty} N_k = 2N.$$

**Solution:** The stated conclusion is false:  $f$  has infinitely many roots. Presumably the intent was to say “roots in  $[0, 1)$ .” Does anyone proofread the Putnam problems before they are printed?

Say the roots of  $f$  in  $[0, 1)$  are  $r_1 < r_2 < \cdots < r_n$  with multiplicities  $m_1, m_2, \dots, m_n$ . Then  $f'$  has a root at  $r_i$  with multiplicity  $m_i - 1$  if  $m_i \geq 2$ ; a root in  $(r_i, r_{i+1})$  for  $1 \leq i \leq n - 1$ ; a root in  $(r_n, 1 + r_1)$ ; and possibly more roots. Thus there are at least  $1 + (n - 1) + \sum_i (m_i - 1) = \sum_i m_i$  roots of  $f'$  in  $[r_1, 1 + r_1)$ , hence in  $[0, 1)$ ; and there are exactly  $\sum_i m_i$  roots of  $f$  in  $[0, 1)$ . Thus  $N_0 \leq N_1$ . By the same argument  $N_1 \leq N_2$ ,  $N_2 \leq N_3$ , etc.

Find  $k_0$  such that  $\sum_{1 \leq j < N} (j/N)^k |a_j/a_N| < 1/2$  for all  $k \geq k_0$ . Abbreviate  $d/dt$  as  $D$ . I will show that  $D^k f$  has exactly  $2N$  roots in  $[0, 1)$  for  $k \geq k_0$ .

Find a real number  $s$  with  $(D^k \sin)(2\pi Ns) = 1$ . Then  $(D^k \sin)(2\pi Nt)$  decreases from 1 at  $s$  to  $-1$  at  $s + 1/2N$ , increases to 1 at  $s + 2/2N$ , etc. By construction

$$\frac{(D^k f)(t)}{(2\pi N)^k a_N} = (D^k \sin)(2\pi Nt) + \sum_{1 \leq j < N} \left(\frac{j}{N}\right)^k \frac{a_j}{a_N} (D^k \sin)(2\pi jt),$$

so  $(D^k f)(t)$  has the same sign as  $a_N (D^k \sin)(2\pi Nt)$  whenever  $|(D^k \sin)(2\pi Nt)| > 1/2$ : in particular, at  $s, s + 1/2N, s + 2/2N, \dots$ . Therefore  $D^k f$  has at least one root in  $[s, s + 1/2N)$ .

It is not possible for  $D^k f$  to have two roots in  $[s, s + 1/2N)$ . Indeed, the roots are in the subinterval  $[s + 1/6N, s + 1/3N]$  where  $(D^k \sin)(2\pi Nt)$  is in  $[-1/2, 1/2]$ . If there were two roots then  $D^{k+1} f$  would also have a root in the subinterval, so  $(D^{k+1} \sin)(2\pi Nt)$  would be in  $[-1/2, 1/2]$ ; contradiction.

The same comments apply to  $[s + 1/2N, s + 2/2N)$  and so on. Thus  $D^k f$  has exactly  $2N$  roots in  $[s, s + 1)$ , hence in  $[0, 1)$ .

### Problem B4

Let  $f(x)$  be a continuous function such that  $f(2x^2 - 1) = 2xf(x)$  for all  $x$ . Show that  $f(x) = 0$  for  $-1 \leq x \leq 1$ .

**Solution:** Thanks to Kahan for pointing out the role of  $\cos$  here. My original solution constructed  $\cos$  manually.

Define  $g(y) = f(\cos 2\pi y)$ . Then  $g$  is continuous;  $g$  is even;  $g$  has period 1; and  $g(2y) = f(\cos 4\pi y) = f(2(\cos 2\pi y)^2 - 1) = 2(\cos 2\pi y)f(\cos 2\pi y) = 2(\cos 2\pi y)g(y)$ .

In particular,  $g(1/3) = g(-1/3) = g(2/3) = -g(1/3)$ , so  $g(1/3) = 0$ . Thus  $g(n+1/3) = 0$  for all integers  $n$ . In fact,  $g((n+1/3)/2^k) = 0$  for all  $n$  and all  $k \geq 0$ . Indeed, if  $k \geq 1$ , then  $g((n+1/3)/2^{k-1}) = 0$  by induction, and  $\cos(2\pi(n+1/3)/2^k) \neq 0$ , so  $g((n+1/3)/2^k) = 0$ .

The set  $\{(n+1/3)/2^k\}$  is dense, so  $g$  is 0 everywhere. Thus  $f$  is 0 on the range of  $\cos$ , namely  $[-1, 1]$ .

Robin Chapman comments that one can remove the  $2 \cos 2\pi y$  factor by considering  $f(\cos 2\pi y)/\sin 2\pi y$  for all non-integer  $y$ .

### Problem B5

Let  $S_0$  be a finite set of positive integers. We define finite sets  $S_1, S_2, \dots$  of positive integers as follows:

Integer  $a$  is in  $S_{n+1}$  if and only if exactly one of  $a-1$  or  $a$  is in  $S_n$ .

Show that there exist infinitely many integers  $N$  for which  $S_N = S_0 \cup \{N+a : a \in S_0\}$ .

**Solution:** Define a polynomial  $f_n$  as  $\sum_{a \in S_n} x^a$ . Then  $f_{n+1} \equiv (x+1)f_n \pmod{2}$ , so  $f_n \equiv (x+1)^n f_0$ .

In particular, if  $n$  is a power of 2 larger than  $\deg f_0$ , then  $f_n \equiv (x+1)^n f_0 \equiv (x^n+1)f_0 = x^n f_0 + f_0$ , and all coefficients of  $x^n f_0 + f_0$  are 0 or 1, so  $f_n = x^n f_0 + f_0$ ; i.e.,  $a \in S_n$  if and only if  $a \in S_0$  or  $a-n \in S_0$ .

### Problem B6

Let  $B$  be a set of more than  $2^{n+1}/n$  distinct points with coordinates of the form  $(\pm 1, \pm 1, \dots, \pm 1)$  in  $n$ -dimensional space, with  $n \geq 3$ . Show that there are three distinct points in  $B$  which are the vertices of an equilateral triangle.

**Solution:** The following solution is a composite of solutions from several other people.

Define  $A = \{(\pm 1, \pm 1, \dots, \pm 1)\}$ . For each  $p \in A$  define  $\Delta_p = \{q \in B : |p-q| = 2\}$ . Then  $\sum_{p \in A} \#\Delta_p = \sum_{q \in B} \#\{p \in A : |p-q| = 2\} = \sum_{q \in B} n = n\#B > 2^{n+1} = 2\#A$ . Thus  $\#\Delta_p > 2$  for some  $p \in A$ . Any distinct  $q_1, q_2, q_3 \in \Delta_p$  form an equilateral triangle in  $B$ .